

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁴ : C07K 3/08, 13/00, 15/00 A61K 35/14, 37/02</p>	<p>A1</p>	<p>(11) International Publication Number: WO 89/ 01486 (43) International Publication Date: 23 February 1989 (23.02.89)</p>
<p>(21) International Application Number: PCT/US88/02700 (22) International Filing Date: 9 August 1988 (09.08.88) (31) Priority Application Numbers: 084,335 228,035 (32) Priority Dates: 11 August 1987 (11.08.87) 5 August 1988 (05.08.88) (33) Priority Country: US (71) Applicant: NEW YORK UNIVERSITY [US/US]; 70 Washington Square South, New York, NY 10012 (US). (72) Inventors: ELSBACH, Peter ; 440 East 23rd Street, New York, NY 10010 (US). WEISS, Jerrold ; 283 Avenue C, New York, NY 10009 (US).</p>		<p>(74) Agents: FRANKFORT, Howard, M. et al.; Darby & Darby, 405 Lexington Avenue, New York, NY 10174 (US). (81) Designated States: AT (European patent), BE (Euro- pean patent), CH (European patent), DE (European patent), FR (European patent), GB (European pa- tent), IT (European patent), JP, LU (European pa- tent), NL (European patent), SE (European patent). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt</i> <i>of amendments.</i></p>
<p>(54) Title: BIOLOGICALLY ACTIVE BACTERICIDAL/PERMEABILITY-INCREASING PROTEIN FRAGMENTS</p>		
<p>(57) Abstract</p> <p>A biologically active polypeptide fragment of mammalian bactericidal/permeability increasing proteins is provided. Also provided are methods for isolating the fragments and methods for employing the polypeptide fragments as antimicrobial agents active against susceptible gram-negative bacteria. Also provided is a purified, isolated DNA sequence encoding human bactericidal/permeability-increasing protein and biologically-active fragments thereof.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	ML	Mali
AU	Australia	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	IT	Italy	NO	Norway
BJ	Benin	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	LI	Liechtenstein	SN	Senegal
CH	Switzerland	LK	Sri Lanka	SU	Soviet Union
CM	Cameroon	LU	Luxembourg	TD	Chad
DE	Germany, Federal Republic of	MC	Monaco	TG	Togo
DK	Denmark	MG	Madagascar	US	United States of America
FI	Finland				

5

10

BIOLOGICALLY ACTIVE BACTERICIDAL/
PERMEABILITY-INCREASING PROTEIN FRAGMENTS

15 BACKGROUND OF THE INVENTION

This application is a continuation-in-part of copending, commonly assigned U.S. patent application Serial No. 084,335, filed August 6, 1987 of Peter Elsbach and Jerrold Weiss.

20 The United States government has rights to this invention by virtue of research grant Nos. DK-05472 and AI-18571 from the National Institute of Health.

The present invention pertains to biologically active, polypeptide fragments of mammalian bactericidal/permeability-increasing proteins and methods for making and using said
25 fragments.

Bactericidal/permeability-increasing protein (BPI) is a 50 to 60 Kd protein, isolated from the granules of mammalian polymorphonuclear leukocytes (PMN) which are blood cells that
30 are essential in the defense against invading microorganisms in mammals. BPI occurs only in cells of the myeloid series of blood cells, is produced at the promyelocytic/myelocytic stage of differentiation and is located in the primary granules in these cells.

35 BPI is a potent bactericidal agent active against a broad range of gram-negative bacterial species. It exhibits a high degree of specificity in its cytotoxic effect, i.e. 10-

40nM (0.5-2.0 micrograms), producing greater than 90% killing of 10^7 sensitive bacteria whereas 100-fold higher concentrations of BPI are non-toxic for other microorganisms and eukaryotic cells. All available evidence suggests that in the
5 intact PMN and in crude leukocyte fractions, BPI is the principal oxygen-independent agent present which is active against BPI-sensitive bacteria.

BPI isolated from both human and rabbit PMN has been purified to homogeneity. The molecular weight of human BPI is
10 approximately 58,000 Daltons (58 kDa) and that of rabbit BPI is approximately 50 kDa. The amino acid composition of these two proteins is closely similar as is the amino acid sequence of their first 15 NH₂-terminal amino acid residues. Both proteins are highly basic, having an isoelectric point greater than 9.6.

15 The biological effects of BPI require attachment of the protein to the surface of the susceptible gram-negative bacteria. Initial binding of BPI to target cells involves electrostatic interactions between the basic protein and the negatively charged sites on the lipopolysaccharides (LPS) on
20 the bacterial outer membrane and leads to an activation of bacterial enzymes that degrade phospholipids and peptidoglycans. The final stage of action is the actual killing of the bacteria by an as yet unknown mechanism. The closely similar amino acid composition and nearly identical bactericidal and membrane-perturbing properties of BPI purified
25 from human and rabbit PMN suggest that this protein has been highly conserved during evolution and is an important member of the anti-bacterial arsenal of the mammalian PMN.

Due to its potent bactericidal action against gram-
30 negative bacteria and lack of cytotoxicity towards other microorganisms and eukaryotic cells, it is envisioned that BPI may be employed as a chemotherapeutic agent and/or as a model for the design of new antibiotic agents. However, due to its large molecular weight (58 kDa for the human holoprotein), both
35 sequencing and determination of the structural organization of BPI have been hampered (hereinafter the entire BPI molecule is referred to as the holoprotein). The possibility has been

raised that, as in the case with other cytotoxic proteins, BPI has a structural organization where the different functions, namely binding, envelope-altering and killing reside in different domains within the BPI molecule. Although BPI fragments, obtained by digestion of the holoproteins with the proteolytic enzyme elastase, has been disclosed (Weiss, J. et al., Clin. Res 34: 537A, 1986), the fragments tested remained associated under the non-denaturing conditions employed. No biological activity was ascribed to any isolated fragments. Moreover, antibodies directed against the holoprotein did not recognize these fragments under denaturing conditions when analyzed using the well-known Western blotting procedure.

Therefore, in light of the above, there is a need in the art for biologically active peptide fragments of BPI for use as bactericidal/permeability increasing agents as well as therapeutic agents. Such fragments are also needed to provide sequence information on BPI to direct the design of future generations of antimicrobial agents specific for gram-negative bacteria and to be used as probes into the molecular organization of the holoproteins.

OBJECTS OF THE INVENTION

It is an object of the present invention to provide biologically active peptide fragments of mammalian BPI.

Another object of the present invention is to provide biologically active peptide fragments of mammalian BPI with improved antimicrobial effectiveness.

Yet another object of the present invention is to provide a process for the production of biologically active peptide fragments of mammalian BPI.

Yet another object of the present invention is to provide methods for treating mammals suffering from infections caused by gram-negative bacteria.

A further object of the present invention is to provide a method for increasing the permeability of gram-negative bacteria.

A still further object of the present invention is to

increase the effectiveness of gram-negative bactericidal agents.

These and other objects of the present invention will be apparent to those of ordinary skill in the art in light of
5 the present description, accompanying claims and appended drawings.

SUMMARY OF THE INVENTION

The present inventors have unexpectedly discovered
10 biologically active fragments of mammalian BPI substantially shorter in length than the native BPI protein. Although these fragments are substantially smaller than the native molecule, they retain at least substantially all of the bactericidal and permeability-increasing properties of the intact protein.

15 The biologically active BPI fragments of the present invention can be produced by incubating a sample comprising the BPI holoprotein under BPI cleaving conditions and recovering biologically active fragments of the BPI holoprotein. The preferred BPI cleaving conditions include heating of the BPI
20 holoprotein in an acceptable buffer for a time ranging between about 16 and 24 hours at a temperature between about 20°C and about 37°C..

In another aspect, the present invention provides a method for treating mammals suffering from infections caused by
25 gram-negative bacteria comprising administering to mammals in need of such treatment a gram-negative bactericidal-effective amount of at least one of the above-mentioned biologically active BPI fragments.

In another aspect, the present invention provides
30 pharmaceutical formulations for treating infections in mammals caused by gram negative bacteria comprising a gram-negative bactericidal-effective amount of at least one of the biologically active BPI fragments or pharmaceutically-acceptable salts thereof.

35 In yet another aspect, the present invention provides a method for increasing the permeability of gram-negative bacteria comprising incubating the bacteria with a gram-

negative-bacterial-permeability-increasing-effective amount of a biologically active fragment of BPI.

In still another aspect, the present invention provides a method for increasing the effectiveness of gram-negative bactericidal agents in mammals in need of such treatments, comprising co-administration of such agents with the biologically active fragments of BPI.

A still further aspect of the present invention is directed to a purified, isolated DNA sequence having the sequence set out in Figure 5 or its complementary strand and DNA sequences hybridizing under stringent hybridization conditions with said DNA sequences. The DNA sequence encodes the human bactericidal/permeability-increasing protein.

15 BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a photograph of a stained SDS-PAGE gel showing the production and purification of the human BPI fragment of a preferred embodiment of the present invention.

Figure 2 is a graph showing the chromatographic behavior of the human BPI holoprotein (a) and human 25 kDa BPI fragment of the present invention (b) on reverse phase HPLC.

Figure 3 is a series of graphs comparing the biological activities of the 25 kDa human BPI fragment of the present invention and the holoprotein toward E. coli J5. (A) bactericidal activity; (B) effect on bacterial protein synthesis; (C) permeability increasing activity; and (D) phospholipase activation.

Figure 4 is a graph comparing the bactericidal effect of the 25 kDa human BPI fragment of the present invention and the holoprotein on E.coli 0111:B4.

Figure 5, bottom line, shows the sequence of the cDNA encoding human BPI whereas printed above is the corresponding amino acid sequence. The two potential glycosylation sites are overlined.

Figure 6 is an autoradiogram of a Northern blot analysis of human BPI mRNA.

Figure 7 is an autoradiogram of a Southern blot

analysis of human genomic DNA using a BPI cDNA probe.

DETAILED DESCRIPTION OF THE INVENTION

The present inventors have unexpectedly discovered
5 biologically active fragments of BPI isolated from mammalian
PMN. NH₂-terminal amino acid sequence analysis demonstrated
that, in the case of human BPI, the fragment represents a
portion of the BPI molecule proximal to the NH₂-terminus, as
shown in Example 3 below. The fragment possesses all of the
10 antibacterial and membrane permeability-increasing functions
contained in the complete molecule but is substantially smaller
(i.e. has a lower molecular weight) than the holoprotein.
"Substantially smaller" is defined herein as being up to about
half the size of the holoprotein. This is a most surprising
15 finding because, in the case of other cellular toxins and
proteins, the entire molecule is necessary for the full
manifestation of their biological effects. For example,
studies on a wide variety of bacterial and plant cytotoxins,
such as diphtheria toxin, cholera toxin and ricin (toxins which
20 do not demonstrate the unique specificity of the BPI holo-
protein) have revealed that individual functions, such as
binding or catalytic activity, can be expressed by isolated
fragments, but that cytotoxicity (comprising both binding to a
cell membrane and intracellular toxic activity) requires
25 essentially the entire molecule.

The BPI fragments of the present invention are as
potent as the holoprotein against rough E.coli, more potent
than the holoprotein against the generally more resistant,
smooth E.coli (on a molar basis), and retain the specificity of
30 the holoprotein towards gram-negative bacteria. This is a par-
ticularly important finding because smooth gram-negative
bacteria (smoothness being due to the presence of longer LPS
chains in the bacterial cell membrane) generally are more
pathogenic than their corresponding rough counterparts.

35 The distinctive size, chromatographic behavior (Figures
1 and 2 below) amino acid content (Table 1 below) and potency
(see Example 4 below) firmly establish that the BPI fragment of

the present invention is a molecular entity distinct from the holoprotein.

Non-limiting examples of the BPI fragments of the present invention are approximately 25 kDa for human and rabbit BPI. The human 25kDa fragment of the preferred embodiment of the present invention was initially isolated after long-term storage (e.g. two months) of the purified holoprotein in a weakly acidic buffer (10mM ammonium acetate, pH4.6) and can be thus generated. However, it is preferable to produce the BPI fragments of the present invention by incubating the holoproteins in an acceptable buffer i.e., a buffer having sufficient buffering capacity at concentrations between about 10 and about 150 mM at a pH ranging between about 6.0 and about 8.0, such as Tris/HCl, phosphate, and preferably HEPES/NaOH (Sigma Chemicals, St. Louis, MO), or mixtures thereof. The preferred pH is 7.4. The incubations may be performed for a period of time broadly ranging between about 16 and 24 hours and preferably 18 hours, at a temperature ranging between about 20°C and about 37°C and preferably 37°C. A particularly preferred condition comprises incubation in 0.1M HEPES/NaOH buffer, pH 7.4 for 18 hours at 37°C. This has led to the conversion of about 50% of the holoprotein into the biologically active fragments of the present invention. Reincubation of the recovered holoprotein, under these conditions again result in formation of the 25 kDa fragment.

The BPI holoproteins, used as starting materials for the production of the biologically active fragments of the present invention, can be obtained from mammalian cells of the myeloid series of blood cells, such as PMN. Although the fragments of the present invention are not limited to a particular mammalian species, it is preferable to employ such fragments isolated from a homologous mammalian species when treating bacterial infections caused by gram-negative bacteria.

In addition, the BPI holoprotein and/or the biologically active fragments of the present invention may be obtained using recombinant DNA techniques employing the sequence information presented below in Example 3 to synthesize DNA

probes for the detection of DNA sequences coding for BPI in complementary DNA or genomic libraries using methods well-known in the art. The gene coding for the BPI holoprotein, or a portion of the gene coding for the 25 kDa fragment of the present invention (and possibly smaller biologically active fragments thereof) may be inserted into a suitable expression vector for the production of biologically active polypeptides.

In one embodiment, human BPI holoprotein can be obtained from PMN isolated from normal blood or from blood from patients with chronic myelocytic leukemia, as detailed in Example 1 below. Alternatively, human BPI can be extracted from the human leukemic cell line HL-60 (available as ATCC CCL 240, American Type Culture Collection, Rockville, MD). The latter have been found to contain approximately 10 micrograms of BPI holoprotein per 10^8 cells. Mature PMN of either normal or leukemic origin contain approximately 60 micrograms per 10^8 cells of the BPI holoprotein and are therefore the preferred starting material.

Once obtained, the mammalian PMN can be fractionated using, for example, the procedures detailed below in Example 1 in order to obtain primary granules (or alternatively by extraction of whole cells with 0.16N sulfuric acid, as described in Elsbach, P. et al., J. Biol. Chem. 254:11000, 1979, incorporated by reference). Such primary granules isolated from PMN or leukemic cell lines contain the bulk of BPI holoprotein activity. The BPI holoprotein can then be extracted and purified using any technique known in the art which yields a biologically active BPI holoprotein. Although crude extracts obtained from such primary granules can be employed as starting materials for the production of the BPI fragments of the present invention, it is preferable to purify the holoprotein before generating the fragments. Preferred extraction and purification techniques for human and rabbit BPI holoproteins are described in Example 1 below.

The amounts of starting, purified, BPI holoprotein to be employed in practicing the present invention preferably should be at least 200 micrograms of purified holoprotein.

Although it is possible to use smaller amounts of material, this may hamper the recovery of the biologically active fragments due to non-specific losses, as is true with many other biologically-active proteins, such as interferons.

5 Although not wishing to be bound by any theory of operation of the present invention, it is believed that the cleavage of the holoprotein to generate the biologically active fragments of the present invention is due to the presence of serine proteases.

10 The protein cleaving conditions necessary for the production of the biologically active BPI fragments of the present invention are broadly within the pH, temperature and time optima of such serine proteases, i.e. pH6.0 - pH 8.0, 20°C - 37°C, 16-24 hours. Such incubation of the BPI holoprotein
15 will produce cleavage at about 25 kDa from the NH₂-terminus of the holoproteins.

 The biologically active BPI fragments of the present invention can be utilized for the treatment of mammals suffering from diseases caused by gram-negative bacteria such as,
20 bacteremia or sepsis. Due to its exquisite selectivity and lack of cytotoxicity toward cells other than gram-negative bacteria, the BPI fragments of the present invention would be particularly useful as specific therapeutic agents. Currently gram-negative bacterial infections, such as those caused by Es-
25 cherichia coli, various species of Salmonella, Klebsiella or Pseudomonas are treated with antibiotics, such as penicillin derivatives, aminoglycosides and chloramphenicol. The effectiveness of antibiotics is limited due to the fact that gram-negative bacilli tend to display significant intrinsic resistance
30 tance to many currently available antibiotics and to readily develop further resistance due to the acquisition of resistance factor plasmids. Under appropriate selective conditions, rapid dissemination of multiple antibiotic resistance among a wide variety of gram-negative pathogens is known to occur.

35 When employed to treat bacteremia (i.e. the presence of bacteria in the blood stream) or sepsis (bacterial contamination of bodily fluids) caused by gram-negative bacteria, the

BPI fragments of the present invention are preferably administered parenterally, and most preferably intravenously in amounts broadly ranging between about 1 microgram and 1000 micrograms and preferably between 10 and about 250 micrograms per treatment. The duration and number of treatments may vary from individual to individual, depending upon the severity of the illness. A typical treatment regime may comprise intravenous administration of about 100 micrograms of the BPI fragments three times a day. To help avoid rapid inactivation of the BPI fragments of the present invention (and indeed the holoproteins) which has been observed in vitro after incubation with serum, the BPI fragments may be coupled with physiologically-acceptable carriers, such as normally occurring serum proteins (e.g. albumin or lysozyme). The BPI fragments of the present invention could also be employed topically to treat mammals suffering from skin infections caused by susceptible gram-negative bacteria which occur in bedridden patients suffering from decubitus ulcers (bed sores) or in burn patients. When employed as a topical antibacterial agent, the BPI fragments may be administered in the same dosages and frequency as described for parenteral administration above.

The BPI fragments of the present invention can be incorporated in pharmaceutical formulations to be used to treat mammals suffering from gram-negative bacterial infections. Pharmaceutical formulations comprising the BPI fragments of the present invention (or physiologically-acceptable salts thereof) as at least one of the active ingredients, would in addition comprise pharmaceutically-acceptable carriers, diluents, fillers, salts and other materials well-known in the art depending upon the dosage form utilized. For example, preferred parenteral dosage forms would comprise a sterile isotonic saline solution, and may comprise between about 1 microgram and 1000 micrograms of the BPI fragments of the present invention covalently coupled to suitable physiologically-acceptable carriers, such as normally occurring serum proteins, for example lysozyme or albumin, to prevent their inactivation. For use in treating mammals with gram-negative

bacterial infections in body fluids largely devoid of (lipo) proteins, such as urine, pharmaceutical formulations may comprise the above amounts of BPI fragments of the present invention and sterile, isotonic saline solutions for irrigation
5 of the urinary tract.

In another preferred embodiment, the BPI fragments of the present invention in amounts ranging between 1 microgram and 1000 micrograms per dose, may be mixed with antibiotics and may be formulated in the same type of preparations used in
10 antibiotic creams (such as Silvadene, Marion Laboratories, Kansas City, MO, Terramycin, Pfipharmecs, New York, New York or Achromycin, Lederle Laboratories, Pearle River, New York) well-known in the art for topical administration.

In another preferred embodiment of the present invention, pharmaceutical formulations for treating mammals suffering from gram-negative bacterial infections may contain the BPI fragments of the present invention in addition to standard amounts (well-known in the art) of antibiotics such as Penicillin-G (available from E.R. Squibb and Sons, Inc., Princeton,
20 New Jersey) or cephalosporins (available from Eli Lilly & Co., Indianapolis, IN). In a particularly preferred embodiment, the BPI fragments of the present invention may be mixed with hydrophobic antibiotics, such as rifampicin (available as RIFAMPIN, CIBA Pharmaceutical CO., Summit, New Jersey), and
25 hydrophobic penicillins such as Penicillin-V Benzathine (Lederle Labs, Pearl River, NY). The increased permeability of gram-negative bacteria after BPI treatment is expected to enhance the effectiveness of such antibiotics which cannot easily enter non-permeabilized bacteria.

30 The BPI fragments of the present invention are expected to be ideally-suited for co-treatment using any antibiotic, immune system cells or factors such as T-cells or interleukin-2, cytotoxic agents or the like, effective against gram-negative bacteria. Because of the increased sensitivity to the
35 fragments of the present invention of the more pathogenic, smooth, gram-negative bacteria, the BPI fragments of the present invention are expected to decrease resistance of such

bacteria to such factors. Substantially simultaneous administration of the fragments of the present invention and the antibiotic of choice is preferred.

5 An example of the above-mentioned embodiment is demonstrated in Example 4 below, wherein actinomycin D (which normally cannot enter and affect gram-negative bacteria due to its hydrophobic properties) significantly inhibited RNA and protein synthesis only in BPI-treated E. coli.

10 In addition, the present inventors have isolated the gene encoding the human BPI holoprotein and have identified and sequenced BPI cDNA isolated from human promyelocytic leukemia cells (HL-60). The nucleotide sequence of the cDNA and the corresponding amino acid sequence of the holoprotein are set out in Figure 5.

15 The sequence information contained in Figure 5 can be employed to synthesize the 25 kDa, biologically active fragment of BPI. In this case, a vector can be generated comprising DNA residues 123 to about 759-780 (or amino acid residues 1 to about 210-220) of Figure 5 using techniques well-known in the art. In addition, smaller sub-fragments of the cDNA of Figure 5 can be generated using, for example, limited Bal31 nuclease digestion of the entire cDNA, to probe for the minimum sequences necessary for BPI biological activities mentioned above.

25 Alternatively, the BPI holoprotein can be obtained after synthesis by suitably transfected or transformed eukaryotic (mammalian or yeast) or prokaryotic cells and the biologically-active 25 kDa fragments mentioned above can be obtained using the techniques described in Example 2 below.

30 The present invention is described further below in specific examples which are intended to illustrate it without limiting its scope.

EXAMPLE 1: ISOLATION AND PURIFICATION OF HUMAN BPI

35 Human leukocytes were obtained from heparin-treated (100-200 U.S.P. units/10 ml) peripheral blood collected by venipuncture from healthy donors and patients with chronic myelocytic leukemia.

Populations of human PMN were obtained in two ways.

(1) PMN's were isolated by the dextran-sedimentation procedure, followed by centrifugation in an Isopaque-Ficoll gradient (Pharmacia Fine Chemicals, Piscataway, NJ) as described (Boyum, A.J., J. Clin. Lab. Invest. Suppl. 97: 77-89, 1968, incorporated by reference). The leukocyte-rich plasma from healthy donors was first diluted with isotonic Krebs-ringer phosphate buffer (pH7.4) to a concentration of 10,000 to 20,000 cells/-microliter before layering on the Isopaque-Ficoll mixture. The cells were washed twice in Krebs-ringer phosphate before use.

(2) Alternatively, leukocyte-rich plasma obtained by leukopheresis (using procedures well-known in the art) of 400 ml of venous blood, from a patient with chronic myelocytic leukemia, was sedimented directly at 1000 X g for five minutes yielding 3.5×10^{10} leukocytes, essentially all of which were PMN. These cells were washed twice with Krebs-ringer phosphate before homogenization.

For extraction of the human BPI holoprotein, the PMN were first disrupted in one of two ways: 1) Granule-rich fractions, containing the bulk of the BPI activity, were obtained by homogenization at 0°C of PMN suspended in 0.34 M sucrose (2×10^8 cells/ml), as described in Weiss, J. et al., J. Biol. Chem. 253: 2664-2672, 1978, incorporated by reference, followed by centrifugation at 400 x g for 10 and 20,000 xg for 30 minutes at 4°C. The granule-rich pellet was extracted with approximately 10 volumes of 0.2 M sodium acetate (pH 4.0), overnight at 4°C with continuous stirring. The extract was collected as a supernatant by centrifugation of the extract at 20,000 x g for 30 minutes.

2) Alternatively, PMN ($2-3 \times 10^8$ cells/ml) were disrupted in distilled water at 0°C with a Potter-Elvehjem glass homogenizer and a motor-driven teflon pestle (Kontes; subsidiary of Kimble Div. of Owens, IL) and extracted at 0°C for 30 minutes with 0.16N sulfuric acid to solubilize the BPI holoprotein. After centrifugation at 23,000 x g for 20 minutes at 4°C to sediment insoluble material, the extract was dialyzed against 200 mM sodium acetate/acetic acid buffer (pH 4.0). The

BPI in these extracts was purified by gel filtration chromatography on a molecular sieving column (SEPHADEX G-75, superfine, Pharmacia Fine Chemicals, Piscataway, NJ) at 4°C. The beads were prepared according to the manufacturer's instructions and
5 equilibrated in the 0.2M sodium acetate (pH 4.0). Using this technique, substantially all of the BPI holoprotein activity was eluted as a single peak (fractions 35-39) corresponding to a discrete protein peak (5-6% of the total protein applied) just after the void volume.

10 The chromatographic fraction containing the human BPI holoprotein was subjected to further chromatography on an ion exchange resin (SP-SEPHADEX, Pharmacia Fine Chemicals, Piscataway, NJ). Protein was applied to the column, equilibrated in 0.1N NaCl-0.2M sodium acetate/acetic acid buffer (pH4.6) and
15 eluted with a stepwise gradient of buffered NaCl (0.3, 0.5 and 0.75M). Human BPI holoprotein eluted in the last step.

Purified human BPI holoprotein was then isolated by high performance liquid chromatography (HPLC) on a reverse phase C-4 (Vydac) column (Sota Chromatography, Crompond, NY)
20 using an HPLC system (Model 332, Beckman Instruments, Fullerton, CA). The column used a linear gradient of acetonitrile (0-95% volume/volume, J.T. Baker Chemical Co., Philipsburg, NJ) in 0.1% trifluoroacetic acid (TFA, Pierce Chemical Co., Rockford, IL). Human BPI eluted at about 70% acetonitrile and
25 was dialyzed against approximately 50 volumes of 10mM ammonium acetate/acetic acid buffer (pH4.6). Purified BPI was stored either in 0.2M sodium acetate/acetic acid buffer (pH 4.0) or in 10mM ammonium acetate/acetic acid buffer (pH4.0) at 4°C.

30 EXAMPLE 2: PRODUCTION OF HUMAN BPI FRAGMENTS

Purified human BPI holoprotein was incubated in 0.1M HEPES-NaOH buffer, pH7.4 for 18 hours, and then analyzed by sodium dodecylsulfate polyacrylamide gel electrophoresis (SDS-PAGE) in 12% polyacrylamide gels using the buffer system
35 containing 0.375M Tris/HCl and 0.1% SDS of Laemmli, U.K., Nature 227: 680-685 1970 incorporated by reference. The results are shown in Figure 1. In Figure 1, lanes A-E were

stained using the well-known Coomassie blue technique and lanes F and G were silver stained using a commercial kit (Bio-Rad, Bio-Rad Labs, Richmond, CA).

Upon incubation in 10mM ammonium acetate buffer (pH 4.6) at 4°C for two months, 10 micrograms of the purified human protein (Figure 1, lane A) fragmented into two species of approximately 35 and 25 kDa (Figure 1, lane B). Incubation of purified human BPI holoprotein (10 micrograms) for 24 hours at 37°C in 0.1M HEPES/NaOH buffer, pH 7.4, enhanced the accumulation of the two species, particularly the 25 kDa species with the concomitant loss of the holoprotein (Figure 1, lane C). Reverse phase HPLC of this incubated mixture, performed as described above for the holoprotein, yielded two major protein peaks, one co-eluting with native human BPI holoprotein and the other eluting slightly earlier (Figure 2). Protein from the later peak migrated on SDS-PAGE as a single 60 kDa species and protein from the earlier peak migrated as a single 25 kDa species (Figure 1, lanes D and E respectively). Fragmentation of the human BPI holoprotein and isolation of the 25 kDa fragment could be repeated with the recovered human holoprotein upon repetition of this procedure, confirming that the 25 kDa fragment was human BPI-derived.

In like manner, rabbit BPI holoprotein, purified as in Example 1 above (500ng, Figure 1, lane F) was fragmented after incubation for 18 hours at 37°C in 0.1M HEPES-NaOH (pH 7.4) into a 25 kDa species (Figure 1, lane G).

EXAMPLE 3: NH₂-TERMINAL AMINO ACID COMPOSITION
AND SEQUENCE ANALYSIS OF BPI FRAGMENTS

The human 25 kDa BPI fragment of the present invention was subjected to amino acid analysis, and the results were compared with the amino acid analysis of purified 60 kDa human holoprotein. Amino acid compositions were determined using a Waters Pico-Tag amino acid analyzer (Waters Associates, Milford, NA) as described (Bidlingmyer, B.A. et al., J. Chrom. 336: 93-104 1984 incorporated, by reference). Samples were pretreated in vacuo for 24 hours at 110°C with 5.7N HCl

containing 0.05% phenol. The results are shown in Table 1.

Table 1

Amino Acid Composition of the 25 kDa Fragment and of the Human BPI Holoprotein

	<u>25 kDa Fragment</u>	<u>Human BPI Holoprotein</u>
	(% of total)	
5		
Asx	8.7	9.1
Glx	8.8	8.9
Ser	15.2	8.8
Gly	7.1	6.4
10 His	2.9	2.8
Arg	3.6	3.7
Thr	3.5	4.6
Ala	4.1	6.0
Pro	5.0	8.1
15 Tyr	2.6	2.7
Val	5.8	7.3
Met	2.3	2.4
Ile	7.3	5.0
Leu	7.7	10.3
20 Phe	4.2	6.1
Lys	11.4	7.6

The values shown above represent the mole fraction (%) of each amino acid and are the mean of three independent determinations. "Asx" stands for asparagine and/or aspartic acid and "Glx" stands for glutamine or glutamic acid.

Amino acid analysis showed that the human 25 kDa fragment was enriched in lysine and serine and contained less non-polar residues when compared with the holoprotein (Table 1).

NH₂-terminal sequence analysis of the human BPI fragment of the present invention and of the holoprotein were performed using the well-known sequential Edman degradation technique (Edman, P. Eur. J. Biochem. 1:80-91, 1967, incorporated by reference) using an amino acid sequencer (Beckman, Model 890C, Beckman Instruments Inc., Fullerton, CA) for the holoprotein or a gas phase sequencer (Applied Biosystems, Model 470A, Applied Biosystems, Inc., Foster City, CA) for the

fragment. Phenylthiohydantoin derivatives of amino acids released sequentially by the Edman degradation process were analyzed by reverse-phase HPLC using an 150mm C-18 column for human BPI (IBM Instruments Inc., Willingford, CT) or an ODS column for the fragment of the present invention (Dupont Zorbax ODS column, E.I. Dupont de Nemours, Wilmington, DE). The results are shown in Table 2 below.

Table 2

10	Holoprotein	V N P G V V V R I S Q K G L D Y A S Q Q
	25 Kd Fragment	V N P G V V V R I S Q K G L D Y A S Q Q
	V = Val, N = Asn, P = Pro, G = Gly, R = Arg, I = Ile, Q = Gln, K = Lys, L = Leu, A = Ala, S = Ser	

15 As can be seen from the data in Table 2, the NH₂-terminal amino acid sequence of the human 25 kDa fragment of the present invention and the holoprotein derived from human PMN were identical in the first 20 amino acid residues, indicating that the fragment was the NH₂-terminal portion of
20 the human holoprotein.

EXAMPLE 4: BIOLOGICAL PROPERTIES OF THE BPI FRAGMENT OF THE PRESENT INVENTION

The antibacterial effects of the 25 kDa human BPI fragment of the present invention were compared with the known activities of the holoprotein. E. coli J5 (obtained from Dr. L. Leive, NIH Bethesda, MD) which produces short-chain lipopolysaccharides (LPS) in galactose-free culture medium, were grown overnight and then subcultured at 37°C in triethanolamine-buffered media as described in Simon, E.G. et al., (Proc. Nat. Acad. Sci. USA 51:877, 1964, incorporated by reference).
30 5X10⁶ E. coli J5 were incubated in a volume of 250 microliters with increasing amounts of either the human holoprotein or the human 25 kDa fragment of the present invention. The effects on
35 bacterial viability were determined either by (1) diluting an aliquot (5X10⁵ bacteria) of the incubation mixture into 2 ml of nutrient broth (Difco Laboratories, Detroit, MI) and measuring

bacterial growth (absorbance at 550nm using a standard spectrophotometer after approximately 4 hours at 37°C); or (2) plating diluted samples on nutrient agar and counting bacterial colonies after overnight incubation at 37°C. The results are presented in Figure 3. In Figure 3, open circles represent BPI holoprotein-treated bacteria and closed circles represent bacteria treated with the human 25 kDa fragment of the present invention.

Figure 3A shows that the isolated 25 kDa fragment of the present invention killed E. coli J5, a strain of bacteria highly sensitive to the holoprotein, in a dose-dependent manner. A linear regression analysis of the data presented in Figure 3A further showed that the fragment was about twice as potent as the holoprotein on a mass basis, meaning that it is about equally potent on a molar basis because the fragment is about half the size of the holoprotein (Figure 3A).

Killing of E. coli by mammalian BPI is initially accompanied by discrete alterations of the outer envelope without causing any apparent damage to the bacterial biosynthetic machinery. Figure 3B shows that even at almost fully lethal doses, both the human holoprotein and the human 25 kDa fragment of the present invention caused little inhibition of bacterial protein synthesis. In contrast, both the fragment and the holoprotein caused nearly a complete inhibition of E. coli J5 protein synthesis when administered in the presence of 50 micrograms/ml of the antibiotic actinomycin D (Merck, Sharp and Dohme, St. Louis, MO, Figure 3C). This effect of actinomycin D reflects increased permeability of the outer membrane of the bacteria permitting the entry of the normally impermeant actinomycin D into the cell where it inhibited RNA and, consequently, protein synthesis. The dose-dependence of the permeability-increasing effect of the fragment of the present invention and the holoprotein was the same as that shown for the bactericidal activity above, and demonstrated that in this respect also the fragment was twice as active as the holoprotein, on a mass basis.

In order to compare the effects of the fragment of the

present invention with the holoprotein on bacterial phospholipids, bacteria were prelabeled during growth with (1-¹⁴C)-oleic acid (New England Nuclear, Boston, MA) as described in Elsbach, P. et al., J. Biol Chem. 254: 11000-11009, 1979 incorporated by reference. Incubation mixtures were supplemented with 0.4% bovine serum albumin (W/V) to capture phospholipid breakdown products (¹⁴C-free fatty acids and ¹⁴C-lysocompounds) permitting their separation from unhydrolyzed bacterial ¹⁴C-phospholipids by filtration through a membrane filter (Millipore HAWP, Millipore Corp. Bedford, MA) to measure phospholipid degradation. The results are shown in Figure 3D.

As shown in Figure 3D, the dose-dependent activation of bacterial phospholipid degrading enzymes by the holoprotein was also produced by the 25 kDa fragment of the present invention, again requiring only half the mass of protein for a comparable effect.

The action of the BPI holoprotein on E. coli is hindered by the presence in the bacterial outer membrane of lipopolysaccharides with long polysaccharide chains ("smooth strains"). The effectiveness of the 25 kDa fragment of the present invention towards a smooth E. coli strain (0111:B4) was compared with that of the holoprotein. E. coli 0111:B4 is a smooth strain bearing longer polysaccharide chains than E. coli J5. Bacteria (1×10^6) were incubated in 125 microliter mixtures with increasing amounts of the BPI holoprotein or the 25 kDa fragment of the present invention. Bacterial viability was measured as above and is expressed as percent of viability of bacteria incubated alone (without any additions). The results are shown in Figure 4.

As can be seen in Figure 4, the 25 kDa fragment of the present invention (closed circles) was about five times more potent than the holoprotein (open circles) towards E. coli 0111:B4. The five fold enhancement in activity of the 25 kDa fragment of the present invention with respect to the holoprotein, suggests that the smaller size of the fragment of the present invention is a factor in facilitating access of the fragment to binding sites at the base of the LPS polysaccharide

chain.

In order to determine if the human 25 kDa fragment of the present invention retained the same cytotoxic specificity towards gram-negative bacteria as the holoprotein, the activities of the 25 kDa fragment and the holoprotein toward a gram-positive bacterium, Micrococcus lysodeikticus (obtained from Dr. M. Salton, New York University, New York, New York) were compared. The bacteria were grown in brain heart infusion broth (Difco Laboratories, Detroit, MI) at 37°C. Bacterial viability was measured as above for E. coli.

Neither the human 25 kDa fragment of the present invention (5-10 micrograms) nor the holoprotein (10-20 micrograms) produced any effect on the viability of Micrococcus lysodeikticus, even at doses twenty times greater than those that are fully lethal towards gram-negative E. coli J5.

The data presented above demonstrate that the spectrum and potency of the antibacterial activities of the human 25 kDa BPI fragment of the present invention are at least equal to and sometimes substantially greater than those of the holoprotein. The data indicate that all of the molecular determinants required for BPI cytotoxicity reside within the portion of the BPI molecule included in the fragment of the present invention.

EXAMPLE 5: CLONING OF THE cDNA OF HUMAN BPI AND IDENTIFICATION OF THE AMINO ACID SEQUENCE

Two synthetic oligonucleotides were designed to encode the 33 amino terminal residues of human BPI. The probes BPI-1 (GTCAATCCTGGTGTGTGGTCAGGATCTCTCAGAAGGGCCTGGATTATGCCTCCCA) and BPI-2 (GCAAGGCACAGCTGCCCTGCAGAAGGAGCTGAAGAGGATCAAGATTCCTGACTAT) were each designed to encode half of the partially known human BPI sequence as previously disclosed in Ooi, C.E. et al., (J. Biol. Chem. 262: 14891-14894, 1987). The probes were kinase labeled with ³²P using standard techniques well-known in the art and used to independently screen a human genomic liver library as disclosed in Lawn, R.M. et al. (Cell 15: 1157-1174, 1978). Six clones were identified among 500,000 plaques which hybridized independently with each probe. The hybridizing

region of one of these clones was sequenced and clearly encoded the amino terminal end of human BPI. This sequence was interrupted by an intron or intervening sequence but nevertheless predicted an additional 22 amino acid residues which preceded the next intron.

Based on the gene sequence, a new DNA probe was then synthesized which corresponded exactly to the encoded 55 amino terminal amino acid residues. This probe was used to screen a small cDNA library prepared from human HL-60 cells (available as ATCC CCL 240, American Type Culture Collection, Rockville, MD) induced with dimethylsulfoxide, DMSO. In the library of the 300,000 plaques, 4 clones were isolated which hybridized with the exact probe. DNA from the clones was isolated and the hybridizing regions were sequenced by the dideoxy chain termination technique of Smith, A.J.H. (Meth. Enzym. 65: 560-580, 1980). The sequence of the longest clone is presented in Figure 5.

As shown in Figure 5, the sequence predicts a 31 amino acid signal peptide, followed by a 456 residue mature protein. The amino terminal sequence determined by protein sequencing of human BPI matches the encoded cDNA exactly. Furthermore, the deduced amino acid composition of the encoded protein corresponds closely to the amino acid composition determined for purified human BPI as disclosed in Ooi, C.E. et al., supra. The encoded sequence predicts a protein of 50.6 kD; the estimated molecular size of purified human BPI is approximately 58 kD. This difference in the apparent size may reflect the presence of two potential N-linked glycosylation sites at positions 122 and 249 of the protein (indicated by overlines in Figure 5).

To further demonstrate that this cDNA encoded human BPI, its expression was engineered in mammalian cells. The entire cDNA was subcloned in a mammalian cell expression vector (Wood, W.I. et al., Nature 312: 330-337, 1984), and then transfected into a human kidney cell line. Small amounts of recombinant BPI were transiently produced and characterized by Western Blotting techniques, showing an immunoreactive band

with a mobility identical to that of native human BPI (results not shown).

The natural expression of BPI in various human tissues was then analyzed by Northern Blot hybridization. RNA was prepared from various tissues (Chirgwin, J.M. et al., Biochem. 24: 5294-5299, 1979), passed over oligo-dT-cellulose and electrophoresed through a formaldehyde agarose gel (Dobner, P.R. et al., Proc. Nat. Acad. Sci. USA 78: 2230-2234, 1981). The gel was transferred to nitrocellulose as described (Thomas, P.S., Proc. Nat. Acad. Sci. USA 77: 5201-5205, 1980) and hybridized under stringent conditions with BPI cDNA.

As shown in Figure 6, the BPI cDNA probe hybridized well with mRNA prepared from the spleen of a patient with chronic myelocytic leukemia. The spleen was heavily infiltrated with immature myeloid cells. The size of the hybridizing signal was approximately 2,000 bases in length, suggesting that the cDNA sequence presented in Figure 5 was full length. The BPI probe did not hybridize with mRNA from normal spleen, mature peripheral blood leukocytes, liver, kidney, or brain (Figure 6). This result is in agreement with previous observations on the location of BPI in various cell types and tissues; the presence of BPI has been previously shown to be restricted to cells of the myeloid series. The BPI cDNA was also used as a probe in Southern hybridizations of human genomic DNA. DNA was isolated from human peripheral blood leukocytes, as described in Blin, N. et al. (Nuc. Acids Res. 3: 2303-2308, 1976), digested with restriction endonucleases Eco RI, BamHI and HindIII, and fractionated on a 1% agarose gel. The DNA was transferred to nitrocellulose (as described in Southern, E.M., J. Molec. Biol. 98: 503-517, 1975) and hybridized with a 5' end fragment of the BPI cDNA probe under stringent conditions (as described in Maniatis et al., Molecular Cloning, a laboratory Manual, pp. 387-389, Cold Spring Harbor Laboratories, NY, 1982, incorporated by reference).

As shown in Figure 7, a single hybridizing band was observed in restriction digests using Eco RI and BamHI when the

5' end of the BPI cDNA was utilized as a probe. This suggested that BPI was encoded by a single gene.

The primary structure of the human BPI protein sequence reveals several features which may be critical for its function. As mentioned above, an amino terminal 25 kD fragment contains all of the bactericidal activity of the holoprotein. A clear charge asymmetry can be observed when the amino terminal 25 kD fragment is compared with the holoprotein. The amino terminal end contains 16 more basic than acidic residues (28 lysine/arginine vs. 12 aspartate/glutamate), while the carboxy terminal end is slightly acidic (20 basic vs. 22 acidic residues). The very basic nature of the amino terminal domain may promote an electrostatic interaction of BPI with the negatively charged LPS on the bacterial envelope.

15

PAPER EXAMPLE I: CO-TREATMENT OF GRAM-NEGATIVE BACTERIA WITH THE HUMAN BPI FRAGMENT AND PENICILLINS

The human BPI fragment of the present invention will be used to test the effectiveness of compositions containing the fragments and Penicillin-G or a hydrophobic derivative, Penicillin-V. Both smooth (*E. coli* 0111:B4) and rough (*E. coli* J5) gram-negative bacteria will be seeded and incubated as in Example 3 above with serial two-fold dilutions containing: the human 25 kDa BPI fragment of the present invention (1 microgram - 1000 micrograms) alone, Penicillin-G (3000 - 300,000 units) alone, Penicillin-V Benzathine (3000 - 300,000 units) alone and compositions containing the same concentrations of the above as mixtures, e.g. the BPI fragment plus Penicillin-G and the BPI fragment plus Penicillin-V. Bacterial viability will be monitored as above in Example 3.

It is expected that lower amounts of both of the penicillins will be effective in killing both smooth and rough *E. coli* strains in the presence of the human 25 kDa BPI fragments showing the efficacy of this embodiment of the present invention.

WHAT IS CLAIMED IS:

1 1. A purified, isolated polypeptide fragment having
2 the properties of bactericidal/permeability-increasing
3 holoprotein, said polypeptide fragment having a substantially
4 lower molecular weight and substantially fewer amino acids than
5 said holoprotein.

1 2. A method for producing a polypeptide fragment
2 having mammalian bactericidal/permeability-increasing
3 holoprotein activity, said polypeptide fragment having a
4 substantially lower molecular weight and substantially fewer
5 amino acids than said holoprotein comprising the steps of:
6 incubating a sample comprising mammalian bac-
7 tericidal/permeability-increasing holoprotein under protein
8 cleaving conditions preserving the bactericidal/permeability-
9 increasing activity in said sample; and
10 subsequently recovering a biologically active
11 fragment of said protein having the properties of mammalian
12 bactericidal/permeability holoprotein but having a substantial-
13 ly lower molecular weight and fewer amino acids than said
14 protein.

1 3. A method for treating infections caused by gram-
2 negative bacteria in mammals comprising administering to
3 mammals in need of such treatment an effective amount of a
4 biologically active polypeptide fragment having the properties
5 of bactericidal/permeability-increasing holoprotein, said
6 polypeptide having a substantially lower molecular weight and
7 substantially fewer amino acids than said bactericidal/per-
8 meability-increasing holoproteins.

1 4. A pharmaceutical formulation for treating infec-
2 tions caused by gram-negative bacteria in mammals comprising an
3 effective amount of a biologically active polypeptide fragment
4 and pharmaceutically-acceptable salts thereof, said polypeptide
5 fragment having the properties of mammalian bactericidal/per-
6 meability-increasing holoprotein and having a substantially

7 lower molecular weight and fewer amino acids than said
8 holoprotein.

1 5. The pharmaceutical formulation of claim 4 further
2 comprising a bactericidal-effective amount of an antibiotic
3 selected from the group consisting of rifampicin, penicillin
4 and cephalosporin.

1 6. A method for increasing the permeability of gram-
2 negative bacteria comprising exposing said gram-negative
3 bacteria to an effective amount of a polypeptide fragment
4 having the properties of bactericidal/permeability increasing
5 holoprotein and having a substantially lower molecular weight
6 and substantially fewer amino acids than said protein.

1 7. A method for killing gram-negative bacteria
2 comprising exposing said bacteria to an amount of a polypeptide
3 having the properties of bactericidal/permeability-increasing
4 holoprotein, said polypeptide having a substantially lower
5 molecular weight and substantially fewer amino acids than
6 bactericidal/permeability-increasing holoprotein, and an amount
7 of bactericidal agent selected from the group consisting of
8 gram-negative bacterial antibiotics, immune system cells,
9 immune factors, cytotoxic agents and mixtures thereof, said
10 amounts in combination being bactericidally effective against
11 said bacteria.

1 8. A purified, isolated DNA sequence comprising the
2 sequence as set out in Figure 5 or its complementary strand and
3 DNA sequences which hybridize under stringent hybridization
4 conditions with said DNA sequences.

1 9. A purified, isolated DNA sequence consisting
2 essentially of a DNA sequence encoding a polypeptide having an
3 amino acid sequence sufficiently duplicative of human bac-
4 tericidal/permeability-increasing protein to allow possession
5 of the biological property of increasing the permeability and

6 killing of susceptible gram negative bacteria.

1 10. A purified, isolated DNA sequence which hybridizes
2 under stringent hybridization conditions with DNA sequences
3 that encode natural bactericidal/permeability-increasing
4 holoprotein.

1 11. A purified, isolated DNA sequence having the
2 sequence of from nucleotide 123 to about nucleotide 780 as set
3 out in Figure 5.

1 12. The DNA sequence of claim 11 comprising a biologi-
2 cally-active fragment of bactericidal/permeability-increasing
3 holoprotein.


1 13. A purified, isolated protein having the properties
2 of naturally occurring human bactericidal/permeability-increas-
3 ing protein.

1 14. The purified, isolated protein of claim 13
2 comprising the amino acid sequence as set out in Figure 5.

1 15. A purified, isolated protein comprising the amino
2 acid sequence from amino acid residue 1 to about amino acid
3 residue 220 as set out in Figure 5.

1/9

FIG. 1



SUBSTITUTE SHEET

FIG. 2a









FIG. 2b



6/2

3/9

FIG. 3A*Bacterial Viability***FIG. 3B** *^{14}C -Amino Acid Incorp.
(- Actinomycin D)***FIG. 3D***Phospholipid Hydrolysis***FIG. 3C** *^{14}C -Amino Acid Incorp.
(+ Actinomycin D)*

4/9




FIG. 4

SUBSTITUTE SHEET

FIG. 5

-30
 met arg glu asn met ala arg gly pro cys asn ala pro arg trp val ser
 ATG AGA GAG AAC ATG GCC AGG GGC CCT TGC AAC GCG CCG AGA TGG GTG TCC
 1 CAGGCCTTGAGGTTTGGCAGCTCTGGAGG
 -10
 leu met val leu val ala ile gly thr ala val thr ala ala Val Asn Pro Gly Val Val Arg Ile Ser Gln
 82 CTG ATG GTG CTC GTC GGC ATA GGC ACC GCG GTG ACA GCG GCC GTC AAC CCT GGC GTC GTG GTC AGG ATC TCC CAG
 -20
 Lys Gly Leu Asp Tyr Ala Ser Gln Gln Gly Thr Ala Ala Leu Gln Lys Glu Leu Lys Arg Ile Lys Ile Pro Asp
 157 AAG GGC CTG GAC TAC GCC AGC CAG CAG GGC ACC GCG GCT CTG CAG AAG GAG CTG AAG AGG ATC AAG ATT CCT GAC
 30
 Tyr Ser Asp Ser Phe Lys Ile Lys His Leu Gly Lys Gly His Tyr Ser Phe Tyr Ser Met Asp Ile Arg Glu Phe
 232 TAC TCA GAC AGC TTT AAG ATC AAG CAT CTT GGG AAG GGG CAT TAT AGC TTC TAC AGC ATG GAC ATC CGT GAA TTC
 50
 Gln Leu Pro Ser Ser Gln Ile Ser Met Val Pro Asn Val Gly Leu Lys Phe Ser Ile Ser Asn Ala Asn Ile Lys
 307 CAG CTT CCC AGT TCC CAG ATA AGC ATG GTG CCC AAT GTG GGC CTT AAG TTC TCC ATC AGC AAC GCC AAT ATC AAG
 80
 Ile Ser Gly Lys Trp Lys Ala Gln Lys Arg Phe Leu Lys Met Ser Gly Asn Phe Asp Leu Ser Ile Glu Gly Met
 382 ATC AGC GGG AAA TGG AAG GCA CAA AAG AGA TTA AAA ATG AGC GGC AAT TTT GAC CTG AGC ATA GAA GGC ATG
 100
 Ser Ile Ser Ala Asp Leu Lys Leu Gly Ser Asn Pro Thr Ser Gly Lys Pro Thr Ile Thr Cys Ser Ser Cys Ser
 457 TCC ATT TCG GCT GAT CTG AAG CTG GGC AGT AAC CCC ACG TCA GGC AAG CCC ACC ATC ACC TGC TCC AGC TGC AGC
 120
 Ser His Ile Asn Ser Val His Val His Ile Ser Lys Ser Lys Val Gly Tyr Leu Ile Gln Leu Phe His Lys Lys
 532 AGG CAC ATC AAC AGT GTC CAC GTG CAC ATC TCA AAG AGC AAA GTC GGG TGG CTG ATC CAA CTC TTC CAC AAA AAA
 140
 150
 160

FIG. 5(cont'd)

170 Ile Glu Ser Ala Leu Arg Asn Lys Met Asn Ser Gln Val Cys Glu Lys Val Thr Asn Ser Val Ser Lys Leu
 607 ATT GAG TCT CGG CTT CGA AAC AAG ATG AAC AGC CAG GTC TGC GAG AAA GTG ACC AAT TCT GTA TCC TCC AAG CTG
 180
 190 Gln Pro Tyr Phe Gln Thr Leu Pro Val Met Thr Lys Ile Asp Ser Val Ala Gly Ile Asn Tyr Gly Leu Val Ala
 682 CAA CCT TAT TTC CAG ACT CTG CCA GTA ATG ACC AAA ATA GAT TCT GTG GCT GGA ATC AAC TAT GGT CTG GTG GCA
 200
 210
 220 Pro Pro Ala Thr Thr Ala Glu Thr Leu Asp Val Gln Met Lys Gly Glu Phe Tyr Ser Glu Asn His Asn Pro
 757 CCT CCA GCA ACC ACG GCT GAG ACC CTG GAG ACC CTG GAT GTA CAG ATG AAG GGG GAG TTT TAC AGT GAG AAC CAC CAC AAT CCA
 230
 240 Pro Pro Phe Ala Pro Pro Val Met Glu Phe Pro Ala Ala His Asp Arg Met Val Tyr Leu Gly Leu Ser Asp Tyr
 832 CCT CCC TTT GCT CCA CCA CCA GTG ATG GAG TTT CCC GCT GCC CAT GAC CGC ATG GTA TAC CTG GGC CTC TCA GAC TAC
 250
 260
 270 Phe Phe Asn Thr Ala Gly Leu Val Tyr Gln Glu Ala Gly Val Leu Lys Met Thr Leu Arg Asp Asp Met Ile Pro
 907 TTC TTC AAC ACA GCC GGG CTT GTA TAC TAC CAA GAG GCT GGG GTC TTG AAG ATG ACC CTT AGA GAT GAC ATG ATT CCA
 280
 290 Lys Glu Ser Lys Phe Arg Leu Thr Thr Lys Phe Phe Gly Thr Phe Leu Pro Glu Val Ala Lys Lys Phe Pro Asn
 982 AAG GAG TCC AAA TTT CGA CTG ACA ACC AAG TTC TTT GGA ACC TTC CTA CCT GAG CTG GCC AAG AAG TTT CCC AAC
 300
 310
 320 Met Lys Ile Gln Ile His Val Ser Ala Ser Thr Pro Pro His Leu Ser Val Gln Pro Thr Gly Leu Thr Phe Tyr
 1057 ATG AAG ATA CAG ATC CAT GTC TCA GCC TCC ACC CCG CCA CAC CTG TCT GTG CAG CCC ACC GGC CTT ACC TTC TAC
 330

SUBSTITUTE SHEET

F/G.5(cont'd)

340 Pro Ala Val Asp Val Gln Ala Phe Ala Val Leu Pro Asn Ser Ser Leu Ala Ser Leu Phe Leu Ile Gly Met His
 1132 CCT GCC GTG GAT GTC CAG GCC TTTT GCC GTC CTC CCC AAC TCC TCC CTG GCT TCC CTC TTC CTG ATT GGC ATG CAC

360 Thr Thr Gly Ser Met Glu Val Ser Ala Glu Ser Asn Arg Leu Val Gly Glu Leu Lys Leu Asp Arg Leu Leu Leu
 1207 ACA ACT GGT TCC ATG GAG GTC AGC GCC GAG TCC AAC AGG CTT GTT GGA GAG CTC AAG CTG GAT AGG CTG CTC CTG

380 Glu Leu Lys His Ser Asn Ile Gly Pro Phe Pro Val Glu Leu Leu Gln Asp Ile Met Asn Tyr Ile Val Pro Ile
 1282 GAA CTG AAG CAC TCA AAT ATT GGC CCC TTC CCG GTT GAA TTG CTG CAG GAT ATC ATG AAC TAC ATT GTA CCC ATT

400 Leu Val Leu Pro Arg Val Asn Glu Lys Leu Gln Lys Gly Phe Pro Leu Pro Thr Pro Ala Arg Val Gln Leu Tyr
 1357 CTT GTG CTG CCC AGG GTT AAC GAG AAA CTA CAG AAA GGC TTC CCT CTC CCG ACG CCG GCC AGA GTC CAG CTC TAC

420 Asn Val Val Leu Gln Pro His Gln Asn Phe Leu Leu Phe Gly Ala Asp Val Val Tyr Lys
 1432 AAC GTA GTG CTT CAC CCT CAC CAG AAC TTC CTG CTG TTC GGT GCA GAC GTT GTC TAT AAA TGA AGGCACCAGGGTGCC

440 1511 GGGGGCTGCAGCCGACCTGTTCCCTGATGGGCTGTGGGCGCACCGGCTGCCCTTCCCGAGGAATCCTCTCCAGATCTTAACCAAGAGCCCTTGCAAC

1611 TTCTTCGACTCAGATTTCAGAAATGNTCTAAACACGAGGAACATATTCATTGGAAAGTGCATGGTGTGTATTTTAGGGATATAGAGCTTCTTTCAAGG

1711 GCTAAGGCTGCAGAGATATTTCCCTCCAGGAATCGTGTTCATTTGTAAACCAAGAAATTTCCATTTGTGCTTCATGAAAAAAACTTCTGGTTTTTTTCAT


1811 GTG---poly-A tail

7/9

① ② ③ ④ ⑤ ⑥ ⑦ ⑧ ⑨ ⑩ ⑪ ⑫ ⑬ ⑭ ⑮ ⑯ ⑰ ⑱ ⑲ ⑳ ㉑ ㉒ ㉓ ㉔ ㉕ ㉖ ㉗ ㉘ ㉙ ㉚ ㉛ ㉜ ㉝ ㉞ ㉟ ㊱ ㊲ ㊳ ㊴ ㊵ ㊶ ㊷ ㊸ ㊹ ㊺ ㊻ ㊼ ㊽ ㊾ ㊿

99

FIG. 6



SUBSTITUTE SHEET

9/9




FIG. 7

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US88/02700

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC IPC (4): C07K 3/08, 13/00, 15/00; A61K 35/14, 37/02 U.S 530/324, 350, 380, 402; 514/12, 21; 424/95		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
U.S	530/324, 350, 380, 402 ; 514/12, 21; 424/95	
Documentation Searched other than Minimum Documentation to the extent that such Documents are included in the Fields Searched ⁸		
Chemical Abstract and Biological Abstract Online Computer Search		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X, P	Chemical Abstracts (Columbus Ohio, USA), Volume 107. No. 17, issued 26, October 1987, (Ooi et. al.) "A 25-KDa amino- terminal fragment carries all the antibac- terial activities of the human neutrophil 60-KDa bactericidal/permeability-increasing protein", page 563, column 2, the abstract no. 152677s, J. Biol. Chem. 1987. 262 (31), 14891-4, (Eng.)	1-7, 13-15
X	Chemical Abstracts (Columbus, Ohio, USA), Volume 91, no. 23, issued 03, December 1979, (Elsbach et. al.) "Separation and purification of a potent bactericidal/ permeability-increasing protein and a closely associated phospholipase A ₂ from rabbit polymorphonuclear leukocytes. Ob- servations on their relationship", page 462, column 1, the abstract no. 191112Z, J. Biol. Chem. 1979, 254(21), 11000-9 (Eng).	1, 3-7 13-15
<p>¹⁰ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
26 September 1988	04 JAN 1989	
International Searching Authority	Signature of Authorized Officer	
ISA/US	Christina Chan	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)

Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
X	<u>Chemical abstracts</u> , (Columbus, Ohio, USA), Volume 106, no. 9, issued 02 March 1987 (Forrest et al.), "Kinetics of the generation and action of Chemical mediators in zymosan-induced inflammation of the rabbit peritoneal cavity" page 466. Column 1, the abstract no. 65594y, Br. J. Pharmacol.1986, 89 (4), 719-30 (Eng).	1,3-7 13-15
X	<u>Biological Abstracts</u> (Philadelphia, Pennsylvania USA), Volume 83,no.9 issued 01, May 1987 (Weiss et. al.), "Cellular and subcellular localization of the bactericidal/permeability- increasing protein of neutrophils". AB-496, column 2, the abstract no. 86173, Blood, 1987, 69 (2), 652-659 (Eng).	1,3-7 13-15
Y	<u>Principles of Protein Structure</u> (new york, USA), seventh printing, issued 1985 (Schulz et. al.), "The covalent structure of proteins", pages 47,58, See pages 47,58 in particular.	2

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

V. ☐ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE ¹

This International search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☐ Claim numbers _____, because they relate to subject matter ^{1,2} not required to be searched by this Authority, namely:

2. ☐ Claim numbers _____, because they relate to parts of the International application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out ^{1,3}, specifically:

3. ☐ Claim numbers _____, because they are dependent claims not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☒ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING ²

This International Searching Authority found multiple inventions in this international application as follows:

I. Claims 1-7 and 13-15 drawn to a process, a polypeptide, a formulation and a therapeutic method.

II. Claims 8-12 drawn to a DNA sequence.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the International application.

2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:

3. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers: 1-7 & 13-15

Telephone practice: Mr. Frankfort advised the Examiner without protest that he only wanted the first invention, claims 1-7 and 13-15 Searched

4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remark on Protest

☐ The additional search fees were accompanied by applicant's protest.

☐ No protest accompanied the payment of additional search fees.

Attachment to Form PCT/ISA/210
Part VI.3.

Telephone Approval:

No payment approved by Mr. Frankfort on September 22, 1988 for Group II; Counsel advised that he has no right to protest for any group not paid for and that any protest must be filed no later than 15 days from the date of mailing of the search report. (Form 210).

Reasons for holding lack of unity of invention:

The invention as defined by Group I (claims 1-7 & 13-15) is drawn to a process, a polypeptide, a formulation and a therapeutic method which is classified in class 530 and 514, subclass 350+ and 12+, respectively and which bears no relationship whatsoever to the invention defined by Group II (claims 8-12) which is drawn to a DNA sequence and is classified in class 435, subclass 172.3.

Time Limit For Filing A Protest

Applicant is hereby given 15 days from the mailing date of this Search Report in which to file a protest of the holding of lack of unity of invention. In accordance with PCT Rule 40.2 applicant may protest the holding of lack of unity only with respect to the groups(s) paid for.